

M235T polymorphism in the angiotensinogen gene and cardiovascular disease: An updated meta-analysis of 39 case-control comparisons

ChuanNan Zhai^{1,2}, HongLiang Cong², Hong Zhang², Kai Hou^{1,2}, Ying Zhang², YingYi Zhang²

¹School of Medicine, NanKai University; Tianjin-China

²Department of Cardiology, Tianjin Chest Hospital; Tianjin-China

ABSTRACT

Objective: M235T polymorphism of the angiotensinogen (AGT) gene has been linked with cardiovascular disease (CVD). The aim of this meta-analysis was to investigate whether combined evidence supports this association.

Methods: A systematic search was conducted for studies published up to October 2018 that evaluate the association between AGT M235T polymorphism and risk of CVD. Case-control studies were identified, and the association between AGT M235T polymorphism and CVD risk was assessed using genetic models.

Results: Thirty-nine comparisons from 38 studies were collected, and a meta-analysis and subgroup analysis was performed based on ethnicity. In the overall population (9225 cases and 8406 controls), the occurrence of CVD was found to be associated with AGT M235T polymorphism in both allelic [T vs. M: odds ratio (OR)=1.16] and recessive (TT vs. MT+MM: OR=1.14) models. In subgroup analyses, a significant association was identified between AGT M235T polymorphism and CVD risk in East Asian subgroups in allelic (T vs. M: OR=1.46), homozygous (TT vs. MM: OR=1.78), dominant (MT+TT vs. MM: OR=1.47), and recessive (TT vs. MT+MM: OR=1.68) models, but there was no significant association in Caucasian populations.

Conclusion: Among East Asians, the AGT variant M235T is associated with CVD risk. However, current evidence suggests that there is no such association in the Caucasian population. (*Anatol J Cardiol* 2019; 21: 222-32)

Keywords: angiotensinogen, genetic polymorphism, cardiovascular disease

Introduction

Cardiovascular disease (CVD) is the main cause of death and leads to over 30% of mortality annually worldwide (1). The general risk factors for CVD include smoking, high body mass index, hypertension, lipid metabolism disorders, and diabetes mellitus, among several other factors (2). Emerging evidence has demonstrated that genetic and environmental factors and polymorphisms also play a crucial role in the occurrence and development of CVD (3, 4). The advancement in single-nucleotide polymorphism (SNP) and genome-wide sequencing technologies has led to an increased number of in-depth studies on the genetics of CVD, and a number of candidate genes have been identified, such as those involved in the regulation of lipid me-

tabolism (5), inflammatory cytokines (6), and the renin-angiotensin-aldosterone system (RAAS) (7).

The RAAS plays a critical role in the pathogenesis of coronary heart disease, and previous studies have determined that it is involved in the progression of hypertension and vascular and left ventricular remodeling (8). Much accumulated evidence has indicated that the RAAS is significantly associated with the initiation and progression of coronary atherosclerosis and thrombogenesis (9). In addition, studies involving angiotensin-converting enzyme (ACE) inhibition and angiotensin II receptor blockade have highlighted the vital role of the RAAS, and gene polymorphism of the RAAS may also affect the efficacy of drug (10). Recently, several genetic variants in the RAAS have been found to be significantly associated with

Address for correspondence: HongLiang Cong, MD, Department of Cardiology, Tianjin Chest Hospital; Taierzhuan South Road No. 291, Jinnan District, 300350, Tianjin-China

Phone: 022-88185111 E-mail: hl_cong@126.com

Accepted Date: 14.03.2019 **Available Online Date:** 25.03.2019

©Copyright 2019 by Turkish Society of Cardiology - Available online at www.anatoljcardiol.com
DOI:10.14744/AnatolJCardiol.2019.75282



CVD risk, such as an insertion/deletion polymorphism in the ACE gene, and T175M and M235T polymorphisms in the angiotensinogen (AGT) gene (11-13). AGT is a crucial determinant of angiotensin II levels, which is an important component of the RAAS. Furthermore, polymorphism in the AGT gene may contribute to atherogenesis in the coronary artery and may be related to the development of CVD (14, 15). The M235T polymorphism has been most widely studied; however, several inconsistent results regarding this polymorphism and CVD risk have been reported. Raygan et al. (16), Bonfim-Silva et al. (17), and Isordia-Salas et al. (18) detected positive correlations, whereas Renner et al. (19), Ranjith et al. (20), and Erbas et al. (21) determined that the AGT M235T polymorphism has no significant effect on the development of CVD. Meta-analyses have been performed to resolve these discrepancies; however, these analyses have been compromised by deficiencies in the sample size, and the results have been either inconclusive or only weakly significant (16). Some of the studies have been limited to Asian populations (22, 23), and several of the most recent studies have not been considered. The aim of the present study was to compile case-control research and updated meta-analyses to explore the association between AGT M235T polymorphism and susceptibility for CVD in a range of populations for more accurate assessment.

Methods

Search strategy

A systematic search of MEDLINE, Embase, China National Knowledge Infrastructure, OVID, ScienceDirect, and WanFang databases was performed to identify epidemiological studies on M235T polymorphisms of the AGT gene and CVD that were published up to October 2018. In the literature searches, various combinations of the keywords "angiotensinogen gene," "AGT," "M235T gene," "genetic polymorphism," "variants," or "variations," "coronary heart disease," "coronary artery disease," "cardiovascular disease," "myocardial infarction," "ischemic heart disease," and "coronary stenosis" were used. Only studies published in English or Chinese were included in the study. The references of all full text papers were examined to identify additional relevant studies. Secondary searches of gray literature were not performed. All retrieved articles were organized using reference manager software (Endnote 6).

Inclusion and exclusion criteria

Inclusion criteria were the following: (1) the study evaluated AGT M235T and CVD risk, (2) original research (case-control studies) or AGT M235T genotype frequencies were provided by case-control status, (3) the study had sufficient data to allow the association between AGT M235T and CVD risk, (4) the study included original data, independent of other studies, and (5) the language of the report was in English or

Chinese. Exclusion criteria were the following: (1) overlapping data, (2) missing information (particularly genotype distributions and studies without controls), after having not received the requested information from the corresponding author, and (3) genome scans investigating linkages without detailed genotype frequencies between cases and controls. Two reviewers independently screened the titles and abstracts for eligibility criteria. Thereafter, the reviewers read the full text of the studies that potentially met the inclusion criteria, and the literature was reviewed to determine the final inclusion of data. For each study, the following information was recorded: first author, year of publication, geographical area, ethnicity, number of cases and controls, genotypes for cases and controls, and evidence of Hardy-Weinberg equilibrium in the controls. If the two reviewers disagreed regarding the inclusions of a study, a consensus was reached through additional review and discussion.

Data extraction

The two reviewers extracted data from each study independently, and any discrepancies were resolved. The information extracted from each article in Tables 1 and 2, including first author, year of publication, country of origin, ethnicity of patients, numbers of cases and controls, AGT genotypes, allele distribution of cases and controls, and outcome, was summarized.

Statistical analysis

Data analysis was conducted using STATA 12.0 software (StataCorp, College Station, TX, USA). The association between AGT M235T polymorphism and CVD susceptibility was assessed in the following genetic models: T versus M (allelic), TT versus MM (co-dominant), MT versus MM (co-dominant), MT+TT versus MM (dominant), and MT+MM versus TT (recessive). Inter-study heterogeneity was tested using Q-statistics. The Mantel-Haenszel method for fixed effects and the DerSimonian and Laird method for random effects were used to estimate pooled effects (24). The fixed effects method was used if the result of the Q test was not significant. Otherwise, the pooled odds ratio (OR) and 95% confidence interval (CI), assuming a random effects model, were calculated. Fixed effects assume that genetic factors have similar effects on CVD susceptibility across all studies, and that the observed variations among studies are caused by chance alone (25). The random effects model assumes that different studies may have substantial diversity and assesses both intra- and inter-study variations (26). A recently developed measure, I^2 , was used to quantify the inconsistency among the studies' results for values of $\geq 50\%$, with large heterogeneity among values of $\geq 75\%$ (27). Data are expressed as OR with 95% CI and two-tailed p-values. A p-value < 0.05 was considered statistically significant. Assessment of publication bias was conducted both visually by using a funnel plot and statistically via Begg's funnel plots

and Egger's bias test, which measures the degree of funnel plot asymmetry (28, 29). The Begg's adjusted rank correlation test was used to assess the correlation between test accuracy estimates and their variances. The deviation of Spearman's rho values from zero provides an estimate of funnel plot asymmetry, where positive values indicate a trend toward higher levels of test accuracy in studies with smaller sample sizes. The Egger's bias test detects funnel plot asymmetry by determining whether the intercept deviates significantly from zero in a regression of the standardized effect estimates against their precision. Meta-regression analysis was applied to evaluate the heterogeneity of the studies.

Results

Search results and characteristics of included studies

Initially, 427 potentially relevant articles were obtained; however, after screening the abstracts, most were determined to be irrelevant to our analysis. Of the remaining 51 articles, 13 articles were removed because of an insufficient number of cases or unusable data. Eventually, 23 studies in English (12-21, 30-42) and 15 in Chinese (43-57), including 39 comparisons of the AGT M235T polymorphism that all adopted the observational study design, satisfied the eligibility criteria (Fig. 1). A total of 39 comparisons from the 38 studies of the AGT M235T polymorphism were included in this updated meta-analysis.

The relevant studies included 9225 cases and 8406 controls (Tables 1 and 2). Reference to the "overall population" indicates meta-analysis without ethnic subdivisions. Ethnicity-specific

meta-analysis was categorized by Caucasian, East Asian, and other races (miscellaneous subgroup).

Association of the AGT M235T polymorphism with CVD risk in the overall population

As shown in Figure 2, significant heterogeneity among studies was observed for the overall population ($P_h < 0.10$ or $I^2 \geq 50\%$). Using the random effect models, M235T was found to be associated with an increased risk of CVD in the allelic (T vs. M: OR=1.16, 95% CI=1.05–1.27, $p < 0.001$) and recessive (TT vs. MT+MM: OR=1.14, 95% CI=1.06–1.23, $p < 0.001$) models.

Association of the M235T polymorphism of the AGT gene with CVD risk in subgroups analysis

When analyses were subdivided according to ethnicity, no associations were noted for Caucasians using any of the five genetic models. However, for the East Asian subgroup, M235T was significantly associated with CVD risk in allelic (T vs. M: OR=1.46, 95% CI=1.13–1.90, $p < 0.001$), homozygous (TT vs. MM: OR=1.78, 95% CI=1.18–2.67, $p = 0.01$), dominant (MT+TT vs. MM: OR=1.47, 95% CI=1.05–2.04, $p = 0.02$), and recessive (TT vs. MT+MM: OR=1.68, 95% CI=1.25–2.27, $p < 0.001$) models. In miscellaneous populations, a significant association between M235T and CVD risk was observed in the allelic model (T vs. M: OR=1.21, 95% CI=1.07–1.36, $p < 0.001$), but no association was observed in the other four genetic models. In subgroup analysis, neither moderate nor large heterogeneity was observed among Caucasians, but true heterogeneity was noted among East Asians (T vs. M: $P_h < 0.10$, $I^2 = 83\%$ and TT vs. MT+MM: $P_h < 0.10$, $I^2 = 81\%$) and miscellaneous populations (MT vs. MM: $P_h = 0.05$, $I^2 = 51.1\%$) (Table 3).

Publication bias and sensitivity analysis

Publication bias was not detected in the analyses of the homozygote, heterozygote, or dominant models ($p > 0.05$, for all). However, publication bias was noted in the analyses of the associations between M235T polymorphisms and CVD risk (allelic model: $P_{Egger} = 0.01$, $P_{Begg} = 0.02$ and recessive model: $P_{Egger} = 0.01$) (Table 4). Sensitivity analyses showed that the present meta-analysis was relatively stable and credible (Fig. 3).

Meta-regression

A meta-regression analysis for several potential sources of heterogeneity, including published year, sample size, age, gender, outcome, and ethnic background, was performed. Single covariates were added in the allelic, homozygote, dominant, and recessive models. The results suggest that the East Asian population (allelic model: $p = 0.006$, homozygote model: $p = 0.010$, dominant model: $p = 0.022$, and recessive model: $p = 0.005$) and study size (homozygote model: $p = 0.042$ and recessive model: $p = 0.010$) contributed to the observed heterogeneity across all studies of the association between AGT M235T polymorphisms and CVD susceptibility.




Figure 1. Flow of studies for meta-analysis

Table 1. Characteristics and genotype frequencies for AGT M235T polymorphism in the included studies

Study	Year	ID	Country	Ethnic	Sample size	Genotypes and allele distribution											
						Case/Control		Cases					Controls				
						MM	MT	TT	M	T	MM	MT	TT	M	T		
Tiret et al.	1995	1	France	Caucasian	630	741	229	301	100	759	501	258	372	111	888	594	
Kamitani et al.	1995	2	Japan	East Asian	103	103	6	31	66	43	163	10	41	52	61	145	
Ko et al.	1997	3	China	East Asian	150	338	4	22	124	30	270	4	54	279	62	612	
Chen et al.	1998	4	China	East Asian	57	76	4	13	40	21	93	13	31	32	57	95	
Sheu et al.	1998	5	China	East Asian	102	145	1	26	75	28	176	1	37	107	39	251	
Pastinen et al.	1998	6	Finland	Caucasian	122	122	48	66	37	162	140	53	64	34	170	132	
Frossard et al.	1998	7	UAE	Caucasian	40	61	14	18	8	46	34	16	26	19	58	64	
Gardemann et al.	1999	8	Germany	Caucasian	1058	511	319	582	157	1220	896	385	585	222	1355	1029	
Winkelmann et al.	1999	9	Germany	Caucasian	122	92	38	54	30	130	114	28	53	11	109	75	
Batalla et al.	2000	10	Spain	Caucasian	220	200	69	99	52	237	203	64	96	40	224	176	
Fomicheva et al.	2000	11	Russia	Caucasian	198	152	63	85	50	211	185	43	75	34	161	143	
Olivieri et al.	2001	12	Italy	Caucasian	247	245	63	124	60	250	244	54	76	27	184	130	
Xie et al.	2001	13	China	East Asian	106	86	8	29	69	45	167	11	30	45	52	120	
Fernández-Arcás et al.	2001	14	Spain	Caucasian	212	180	59	121	32	239	185	34	97	49	165	195	
Ermis et al.	2002	15	Turkey	Miscellaneous	102	114	32	48	22	112	92	39	59	16	137	91	
Hooper et al.	2002	16	USA	Miscellaneous	110	185	4	29	67	37	163	2	31	67	35	165	
Zhu et al.	2002	17	China	East Asian	41	116	2	7	32	11	71	18	47	51	83	149	
Zhu et al.	2002	18	China	East Asian	118	106	14	48	56	76	160	10	42	54	62	150	
Bis et al.	2003	19	USA	Caucasian	208	717	71	98	39	240	176	215	349	153	779	655	
Gu et al.	2003	20	China	East Asian	129	90	12	31	86	55	203	7	30	53	44	136	
Ranjith et al.	2004	21	India	Miscellaneous	195	300	24	80	91	128	262	29	127	144	185	415	
Zhu et al.	2004	22	China	East Asian	192	98	12	75	105	99	285	8	36	54	52	144	
Li et al.	2004	23	China	East Asian	120	80	11	60	49	82	158	14	41	25	69	91	
Tobin et al.	2004	24	England	Caucasian	547	505	212	252	83	676	418	197	226	82	620	390	
Ren et al.	2005	25	China	East Asian	100	70	2	10	35	14	80	13	26	31	52	88	
Araujo et al.	2005	26	Brazil	Caucasian	110	104	46	52	12	144	76	43	51	10	137	71	
Renner et al.	2005	27	Austria	Caucasian	1370	733	NA	NA	NA	1537	1203	NA	NA	NA	832	634	
Liang et al.	2006	28	China	East Asian	133	154	2	30	101	34	232	10	60	84	80	228	
Tsai et al.	2007	29	China	East Asian	735	519	15	195	525	225	1245	5	111	403	121	917	
Niu et al.	2008	30	China	East Asian	105	110	8	32	65	48	162	9	47	54	65	155	
Zhu et al.	2010	31	China	East Asian	151	127	9	27	115	45	257	20	51	56	91	163	
Peng et al.	2011	32	China	East Asian	196	200	14	54	128	82	155	18	86	96	122	278	
Konopka et al.	2011	33	Poland	Caucasian	100	95	30	46	24	106	94	22	44	29	88	102	
Mehri et al.	2011	34	Tunisia	Miscellaneous	123	144	29	53	41	111	135	53	61	30	167	121	
Raygan et al.	2016	35	Iran	Miscellaneous	155	185	42	79	34	163	147	71	85	29	227	143	
Bonfim-Silva et al.	2016	36	Brazil	Miscellaneous	153	113	23	69	61	115	191	13	63	37	89	137	
				Caucasian	306	142	73	145	88	291	321	34	68	40	136	148	
Erbas et al.	2017	37	Turkey	Miscellaneous	117	106	11	104	2	126	108	16	85	5	117	95	
Isordia-Salas et al.	2018	38	Mexico	Miscellaneous	242	242	138	98	6	374	110	170	62	10	402	82	

NA - not available




Figure 2. Forest plots of the association between AGT M235T polymorphism and CVD risk. (a) Allelic model, (b) homozygote model, (c) heterozygote model, (d) dominant model, and (e) recessive model

Table 2. Baseline characteristics of the included studies

Study	Ethnic	Outcome	Genotyping-methods	Age		Gender (M/F)		HWE
				Case	Control	Case	Control	
Tiret et al.	C	MI	PCR	53.0±0.3	54.0±0.3	630/0	741/0	Y
Kamitani et al.	EA	MI	PCR	52±1	54±1	103/0	103/0	Y
Ko et al.	EA	MI	PCR	61.5±0.6	56.0±0.6	NR	181/157	Y
Chen et al.	EA	MI	PCR	67.7±8.5	65.7±8.2	50/7	69/7	Y
Sheu et al.	EA	CAD	PCR	NR	NR	102/0	145/0	Y
Pastinen et al.	C	MI	PCR	57.7±4.9	57.7±4.9	122/0	122/0	Y
Frossard et al.	C	MI	PCR	55.0±11.3	53.7±14.0	25/15	31/30	Y
Gardemann et al.	C	MI	PCR	62.2±9.5	58.5±10.6	1058/0	511/0	Y
Winkelmann et al.	C	MI	PCR	55.7±9.6	55.7±9.6	122/0	92/0	Y
Batalla et al.	C	MI	PCR	43±5	42±6	220/0	200/0	Y
Fomicheva et al.	C	MI	PCR	67 (55-85)	11 (6-17)	198/0	152/0	Y
Olivieri et al.	C	MI	PCR	57.7±12.8	59.6±9.5	160/85	221/26	Y
Xie et al.	EA	CAD	PCR	61.4±9.5	52.8±8.7	82/24	54/32	Y
Fernández-Arcás et al.	C	MI	PCR	54±13	56±15	212/0	180/0	Y
Ermis et al.	M	MI	PCR	42.1±11.8	40.3±12.8	NR	NR	Y
Hooper et al.	M	MI	PCR	NR	NR	NR	NR	Y
Zhu et al.	EA	MI	PCR	59.6±10.4	56.6±10.4	27/14	67/49	Y
Zhu et al.	EA	CAD	PCR	NR	NR	NR	NR	Y
Bis et al.	C	MI	PCR	63.6	64.4	128/80	371/346	Y
Gu et al.	EA	CAD	PCR	65.8±9.2	65.3±9.8	81/48	54/36	Y
Ranjith et al.	M	MI	PCR	18-45	18-45	NR	NR	Y
Zhu et al.	EA	CAD	PCR	NR	NR	NR	NR	Y
Li et al.	EA	CAD	PCR	61.5±11.8	59.3±10.5	80/40	47/33	Y
Tobin et al.	C	MI	PCR	61.9±9.2	58.6±10.7	372/175	313/192	Y
Ren et al.	EA	CAD	PCR	60.0±9.8	57.9±11.6	71/29	38/32	Y
Araujo et al.	C	MI	PCR	>18	>18	73/37	44/60	Y
Renner et al.	C	MI	PCR	63.1±10.4	58.4±12.1	1081/289	378/355	NR
Liang et al.	EA	CAD	PCR	64±8	63±8	100/33	116/38	Y
Tsai et al.	EA	CAD	PCR	63.8±11.4	58.6±13.1	531/204	269/250	Y
Niu et al.	EA	CAD	PCR	59±7	57±9	69/36	71/39	Y
Zhu et al.	EA	CAD	PCR	59.7±11.3	58.1±10.8	96/55	71/56	Y
Peng et al.	EA	CAD	PCR	70.0±8.3	69.0±6.4	128/68	132/68	Y
Konopka et al.	C	MI	PCR	57±10	38±11	79/21	76/19	Y
Mehri et al.	M	MI	PCR	62.3±11.8	60.4±10.3	71/52	83/61	Y
Raygan et al.	M	MI	PCR	62.4±3.2	61.7±4.3	102/53	127/58	Y
Bonfim-Silva et al.	M	CAD	PCR	55.7±7.9	51.8±8.4	99/54	49/64	Y
	C	CAD	PCR	55.7±6.7	53.0±7.7	204/102	65/77	Y
Erbas et al.	M	CAD	PCR	50.2±12.3	41.4±11.3	55/62	14/92	Y
Isordia-Salas et al.	M	MI	PCR	41.0±5.3	39.7±5.0	191/51	192/50	Y

C - Caucasian; EA - East Asian; M - Miscellaneous; MI - myocardial infarction; CAD - coronary artery disease; PCR - polymerase chain reaction; NR - no reported; HWE - Hardy-Weinberg equilibrium; Y - yes

Discussion

The AGT gene (on chromosome 1q42–43) comprises five exons and four introns spanning 12 kb, with the M235T polymor-

phism in exon 2. The M235T variant has been demonstrated to alter plasma AGT levels (58, 59), with elevated levels of serum AGT for patients carrying the T allele (60). Furthermore, a positive correlation exists between AGT M235T genotype and

Table 3. Overall and subgroup meta-analysis of the association between AGT M235T polymorphism and risk of cardiovascular disease

Categories	n	T vs. M			TT vs. MM			MT+TT vs. MM			TT vs. MM+MT					
		OR (95% CI)	P	I ² (%) / P _h	OR (95% CI)	P	I ² (%) / P _h	OR (95% CI)	P	I ² (%) / P _h	OR (95% CI)	P	I ² (%) / P _h			
Overall	22	1.16 (1.05-1.27)	0.003	73.3/0.00	1.20 (1.00-1.45)	0.05	56.5/0.00	1.06 (fixed) (0.97-1.15)	0.20	11.7/0.27	1.12 (0.99-1.26)	0.085	42.0/0.00 (1.06-1.23)	1.14 (1.06-1.23)	0.003	73.3/0.00
Subgroup (by population)																
Caucasian		0.99 (fixed) (0.93-1.04)	0.584	34.1/0.10	0.95 (0.79-1.16)	0.63	43.4/0.04	1.00 (fixed) (0.91-1.11)	0.94	00.0/0.54	0.99 (0.89-1.09)	0.761	05.0/0.40 (0.82-1.16)	0.97 (0.82-1.16)	0.743	50.1/0.02
East Asian		1.46 (1.13-1.90)	0.004	83.0/0.00	1.78 (1.18-2.67)	0.01	53.8/0.01	1.05 (fixed) (0.80-1.38)	0.73	00.0/0.84	1.47 (1.05-2.04)	0.023	34.2/0.09 (1.25-2.27)	1.68 (1.25-2.27)	0.001	81.0/0.00
Miscellaneous		1.21 (fixed) (1.07-1.36)	0.002	40.8/0.11	1.22 (0.81-1.84)	0.33	45.0/0.08	1.23 (0.90-1.70)	0.20	51.1/0.05	1.31 (0.98-1.75)	0.065	45.5/0.08 (0.98-1.46)	1.20 (fixed) (0.98-1.46)	0.082	32.8/0.17

n - study numbers, OR - odds ratio, CI - confidence interval, bold values represent statistically significant findings, P_h - P heterogeneity (P<0.1 was considered as a significant difference), fixed - the fixed effects model

Table 4. Publication bias assessment of this meta-analysis

Genetic model	Egger's test		Begg's test	
	t-value	P	t-value	P
Allelic model	2.70	0.01	2.42	0.02
Homozygote model	1.84	0.07	1.28	0.20
Heterozygote model	-0.2	0.85	0.10	0.92
Dominant model	1.18	0.24	0.85	0.39
Recessive model	2.83	0.01	1.91	0.06

P<0.05 was considered as a significant difference

plasma AGT levels in survivors of myocardial infarction (MI) (14). Elevation circulating AGT levels are associated with an increase in the concentration of angiotensin II, which activates cardiomyocyte hypertrophy and fibroblast proliferation by stimulation the AT1 receptor (61, 62). In addition, angiotensin II stimulates vascular apoptosis and may promote the retention of low-density lipoprotein in the coronary arteries, oxidize, and be assimilated by phagocytes, ultimately contributing to the dysfunction of the vascular endothelium and myocardial ischemia and rupture of atherosclerotic plaque (63-65). These processes all play critical roles in promoting the pathological development of CVD.

In previous studies, the distribution of the AGT M235T variant has been shown to differ significantly among various populations. Katsuya et al. (66) demonstrated that homozygous AGT T235 is an independent risk factor, which carries a two-fold increased risk of CVD. In contrast, Tiret et al. (31) suggested that the AGT genetic polymorphism can lead to predisposition to hypertension, with no relationship to CVD. Renjith et al. (20) and Renner et al. (19) reported similar results showing that AGT genotypes are neither related to CVD nor high blood pressure. Batalla et al. (12) and Bonfim-Silva et al. (17) provided evidence of the synergistic effect between AGT polymorphism and CVD, suggesting that M235T polymorphism is significantly correlated with MI and hypertension. Recently, Raygan et al. (16) suggested that M235T polymorphism in Asians can be a useful biomarker for screening of individuals susceptible to MI. Given the disparity of the results, we sought to provide an updated meta-analysis to resolve the discrepancies among studies.

Our results clearly demonstrate a difference in the association of M235T polymorphism among Asians and Caucasians, suggesting that there is heterogeneity based on ethnicity. Our results suggest that the T allele is a genetic risk factor for CVD in Asians, as well as in the miscellaneous population. Previous meta-analyses based on Chinese populations have shown similar positive results (22, 23). Although two recent meta-analyses (16, 67) reported an association between the AGT M235T variant and MI, they did not include recently published studies, and the literature sample size was smaller than the current analysis. Thus, an update of previous meta-analyses that included diverse populations was warranted. Our results include the largest sam-




Figure 3. Sensitivity analyses of the association between AGT M235T polymorphism and CVD risk. (a) Allelic model, (b) homozygote model, (c) heterozygote model, (d) dominant model, and (e) recessive model

ple size to date and provide an ethnicity-based explanation for different results among studies. The association of AGT M235T with stroke was not addressed in the present study but would be an important topic for consideration in future studies to maintain a relatively narrow focus.

The following potential factors may account for differences observed among the various ethnic groups: (1) population diversity (68), (2) different habits among populations (69), and (3) environmental factors leading to differences in susceptibility to CVD (70). Furthermore, we speculate that the non-significant association among Caucasians may be due to the relatively low frequency of the TT genotype in this population. Moreover, owing to the limited number of these studies, the miscellaneous ethnic subgroups were not analyzed further. Therefore, more studies with a larger sample size may reveal factors that influence differences in the association of AGT M235T and CVD, especially among Caucasians and other ethnic populations. The association between AGT M235T and CVD among Asians is evident.

The heterogeneity of associations across all included studies should be noted as it may potentially affect the strength of the present study. Thus, the random effects model was used, and our analysis was based on different ethnic subgroups. Ethnic background and sample size were found to be factors of heterogeneity. However, heterogeneity was still high within the East Asian subgroup. The heterogeneity of results among those included in these studies may be explained by the quality of the included studies, classification of CVD, and sampling criteria. The heterogeneity of genetic effects among individual studies may also be caused by the existence of genetic and environmental or genetic interactions.

Study limitations

The primary limitations of our meta-analysis include: (1) significant publication bias in the allelic and recessive models, (2) insufficient genotyping data of AGT M235T in miscellaneous races, which limited the ability to draw conclusions regarding this population, and (3) potential heterogeneity of clinical variables, such as the general condition of subjects, their medical history, medication compliance, complications of CVD, and other factors.

Conclusion

The genetic polymorphism of AGT M235T is associated with a critical risk of CVD in East Asian populations, with no detectable association in Caucasian populations. However, further studies with multiple ethnicities and rigorous designs should be performed to confirm these conclusions.

Acknowledgment: The authors thank Professors Li and Liu for their great advices on studies of the unpublished literature.

Ethical approval: Ethical approval was not required for this study design.

Conflict of interest: None declared.

Peer-review: Externally peer-reviewed.

Authorship contributions: Concept – C.Z.; Design – C.Z., H.C.; Supervision – C.Z., H.C.; Fundings – H.C.; Materials – C.Z., H.Z., K.H.; Data collection &/or processing – C.Z., H.Z., Y.Z., Y.Y.Z.; Analysis &/or interpretation – C.Z., H.Z., K.H.; Literature search – K.H., Y.Z.; Writing – C.Z., Y.Y.Z.; Critical review – Y.Z., Y.Y.Z.

References

1. Roth GA, Huffman MD, Moran AE, Feigin V, Mensah GA, Naghavi M, et al. Global and regional patterns in cardiovascular mortality from 1990 to 2013. *Circulation* 2015; 132: 1667-78.
2. Ounpuu S, Negassa A, Yusuf S. INTER-HEART: A global study of risk factors for acute myocardial infarction. *Am Heart J* 2001; 141: 711-21.
3. Roberts R, Stewart AF. Genes and coronary artery disease: where are we? *J Am Coll Cardiol* 2012; 60: 1715-21.
4. Meininger JC, Hayman LL, Coates PM, Gallagher PR. Genetic and environmental influences on cardiovascular disease risk factors in adolescents. *Nurs Res* 1998; 47: 11-8.
5. Xia J, Cai W, Peng C. Association of APOA5 T1131C polymorphism and risk of coronary artery disease. *Int J Clin Exp Med* 2015; 8: 8986-94.
6. Yuepeng J, Zhao X, Zhao Y, Li L. Gene polymorphism associated with TNF- α (G308A) IL-6 (C174G) and susceptibility to coronary atherosclerotic heart disease: A meta-analysis. *Medicine (Baltimore)* 2019; 98: e13813.
7. Ryu SK, Cho EY, Park HY, Im EK, Jang YS, Shin GJ, et al. Renin-angiotensin-aldosterone system (RAAS) gene polymorphism as a risk factor of coronary in-stent restenosis. *Yonsei Med J* 2002; 43: 461-72.
8. Wang JG, Staessen JA. Genetic polymorphisms in the renin-angiotensin system: relevance for susceptibility to cardiovascular disease. *Eur J Pharmacol* 2000; 410: 289-302.
9. Koh KK, Han SH, Oh PC, Shin EK, Quon MJ. Combination therapy for treatment or prevention of atherosclerosis: focus on the lipid-RAAS interaction. *Atherosclerosis* 2010; 209: 307-13.
10. Heidari F, Vasudevan R, Mohd ASZ, Ismail P, Etemad A, Pishva SR, et al. Association of insertion/deletion polymorphism of angiotensin-converting enzyme gene among Malay male hypertensive subjects in response to ACE inhibitors. *J Renin Angiotensin Aldosterone Syst* 2015; 16: 872-9.
11. Kondo H, Ninomiya T, Hata J, Hirakawa Y, Yonemoto K, Arima H, et al. Angiotensin I-converting enzyme gene polymorphism enhances the effect of hypercholesterolemia on the risk of coronary heart disease in a general Japanese population: the hisayama study. *J Atheroscler Thromb* 2015; 22: 390-403.
12. Batalla A, Alvarez R, Reguero JR, Hevia S, Iglesias-Cubero G, Alvarez V, et al. Synergistic effect between apolipoprotein E and angiotensinogen gene polymorphisms in the risk for early myocardial infarction. *Clin Chem* 2000; 46: 1910-5.
13. Gardemann A, Stricker J, Humme J, Nguyen QD, Katz N, Philipp M, et al. Angiotensinogen T174M and M235T gene polymorphisms are associated with the extent of coronary atherosclerosis. *Atherosclerosis* 1999; 145: 309-14.
14. Winkelmann BR, Russ AP, Nauck M, Klein B, Böhm BO, Maier V, et al. Angiotensinogen M235T polymorphism is associated with plas-

- ma angiotensinogen and cardiovascular disease. *Am Heart J* 1999; 137: 698-705.
15. Hooper WC, Dowling NF, Wenger NK, Dilley A, Ellingsen D, Evatt BL. Relationship of venous thromboembolism and myocardial infarction with the renin-angiotensin system in African-Americans. *Am J Hematol* 2002; 70: 1-8.
 16. Raygan F, Karimian M1, Rezaeian A, Bahmani B, Behjati M. Angiotensinogen-M235T as a risk factor for myocardial infarction in Asian populations: a genetic association study and a bioinformatics approach. *Croat Med J* 2016; 57: 351-62.
 17. Bonfim-Silva R, Guimarães LO, Souza Santos J, Pereira JF, Leal Barbosa AA, Souza Rios DL. Case-control association study of polymorphisms in the angiotensinogen and angiotensin-converting enzyme genes and coronary artery disease and systemic artery hypertension in African-Brazilians and Caucasian-Brazilians. *J Genet* 2016; 95: 63-9.
 18. Isordia-Salas I, Alvarado-Moreno JA, Jiménez-Alvarado RM, Hernández-Juárez J, Santiago-Germán D, Leños-Miranda A, et al. Association of renin-angiotensin system genes polymorphisms and risk of premature ST elevation myocardial infarction in young Mexican population. *Blood Coagul Fibrinolysis* 2018; 29: 267-74.
 19. Renner W, Nauck M, Winkelmann BR, Hoffmann MM, Scharnagl H, Mayer V, et al. Association of angiotensinogen haplotypes with angiotensinogen levels but not with blood pressure or coronary artery disease: the Ludwigshafen Risk and Cardiovascular Health Study. *J Mol Med (Berl)* 2005; 83: 235-9.
 20. Ranjith N, Pegoraro RJ, Rom L, Lanning PA, Naidoo DP. Renin-angiotensin system and associated gene polymorphisms in myocardial infarction in young South African Indians. *Cardiovasc J S Afr* 2004; 15: 22-6.
 21. Erbas T, Cinar N, Dagdelen S, Gedik A, Yorgun H, Canpolat U, et al. Association between ACE and AGT polymorphism and cardiovascular risk in acromegalic patients. *Pituitary* 2017; 20: 569-77.
 22. Li X, Li Q, Wang Y, Li Y, Ye M, Ren J, et al. AGT gene polymorphisms (M235T, T174M) are associated with coronary heart disease in a Chinese population. *J Renin Angiotensin Aldosterone Syst* 2013; 14: 354-9.
 23. Wang YJ, Pan Y. Angiotensinogen gene M235T polymorphism and risk of coronary artery disease: a meta-analysis. *Mol Med Rep* 2012; 6: 884-8.
 24. Robins J, Greenland S, Breslow NE. A general estimator for the variance of the Mantel-Haenszel odds ratio. *Am J Epidemiol* 1986; 124: 719-23.
 25. Egger M, Smith GD, Phillips AN. Meta-analysis: principles and procedures. *BMJ* 1997; 315: 1533-7.
 26. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7: 177-88.
 27. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; 21: 1539-58.
 28. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; 315: 629-34.
 29. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994; 50: 1088-101.
 30. Kamitani A, Rakugi H, Higaki J, Ohishi M, Shi SJ, Takami S, et al. Enhanced predictability of myocardial infarction in Japanese by combined genotype analysis. *Hypertension* 1995; 25: 950-3.
 31. Tiret L, Ricard S, Poirier O, Arveiler D, Cambou JP, Luc G, et al. Genetic variation at the angiotensinogen locus in relation to high blood pressure and myocardial infarction: the ECTIM Study. *J Hypertens* 1995; 13: 311-7.
 32. Pastinen T, Perola M, Niini P, Terwilliger J, Salomaa V, Vartiainen E, et al. Array-based multiplex analysis of candidate genes reveals two independent and additive genetic risk factors for myocardial infarction in the Finnish population. *Hum Mol Genet* 1998; 7: 1453-62.
 33. Frossard PM, Hill SH, Elshahat YI, Obineche EN, Bokhari AM, Lestringant GG, et al. Associations of angiotensinogen gene mutations with hypertension and myocardial infarction in a gulf population. *Clin Genet* 1998; 54: 285-93.
 34. Fomicheva EV, Gukova SP, Larionova-Vasina VI, Kovalev YR, Schwartz EI. Gene-gene interaction in the RAS system in the predisposition to myocardial infarction in elder population of St. Petersburg (Russia). *Mol Genet Metab* 2000; 69: 76-80.
 35. Fernández-Arcás N, Dieguez-Lucena JL, Muñoz-Morán E, Ruiz-Galdón M, Espinosa-Caliani S, Aranda-Lara P, et al. Both alleles of the M235T polymorphism of the angiotensinogen gene can be a risk factor for myocardial infarction. *Clin Genet* 2001; 60: 52-7.
 36. Olivieri O, Stranieri C, Girelli D, Pizzolo F, Grazioli S, Russo C, et al. Homozygosity for angiotensinogen 235T variant increases the risk of myocardial infarction in patients with multi-vessel coronary artery disease. *J Hypertens* 2001; 19: 879-84.
 37. Ermis C, Tsai MY, Hanson NQ, Akar N, Aras O. Angiotensin I converting enzyme, angiotensin II type 1 receptor and angiotensinogen polymorphisms and early myocardial infarction in Turkish population. *Thromb Haemost* 2002; 88: 693-4.
 38. Bis JC, Smith NL, Psaty BM, Heckbert SR, Edwards KL, Lemaitre RN, et al. Angiotensinogen Met235Thr polymorphism, angiotensin-converting enzyme inhibitor therapy, and the risk of nonfatal stroke or myocardial infarction in hypertensive patients. *Am J Hypertens* 2003; 16: 1011-7.
 39. Tobin MD, Braund PS, Burton PR, Thompson JR, Steeds R, Channer K, et al. Genotypes and haplotypes predisposing to myocardial infarction: a multilocus case-control study. *Eur Heart J* 2004; 25: 459-67.
 40. Araújo MA, Goulart LR, Cordeiro ER, Gatti RR, Menezes BS, Lourenço C, et al. Genotypic interactions of renin-angiotensin system genes in myocardial infarction. *Int J Cardiol* 2005; 103: 27-32.
 41. Mehri S, Mahjoub S, Farhati A, Bousaada R, Ben Arab S, Baudin B, et al. Angiotensinogen gene polymorphism in acute myocardial infarction patients. *J Renin Angiotensin Aldosterone Syst* 2011; 12: 42-7.
 42. Konopka A, Szperl M, Piotrowski W, Roszczyńko M, Stępińska J. Influence of renin-angiotensin system gene polymorphisms on the risk of ST-segment-elevation myocardial infarction and association with coronary artery disease risk factors. *Mol Diagn Ther* 2011; 15: 167-76.
 43. Ko YL, Ko YS, Wang SM, Chu PH, Teng MS, Cheng NJ, et al. Angiotensinogen and angiotensin-I converting enzyme gene polymorphisms and the risk of coronary artery disease in Chinese. *Hum Genet* 1997; 100: 210-4.
 44. Sheu WH, Lee WJ, Jeng CY, Young MS, Ding YA, Chen YT. Angiotensinogen gene polymorphism is associated with insulin resistance in nondiabetic men with or without coronary heart disease. *Am Heart J* 1998; 136: 125-31.
 45. Chen D, Zhang M, Fan W, Shi H, Li Y, Chen Q, et al. A molecular variant of angiotensinogen gene is associated with myocardial infarction in Chinese. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 1998; 15: 133-5.
 46. Xie Y, Zhang Z, Ma W, Yang D, Geng Z, Jia Y, et al. Association between gene polymorphism of angiotensinogen (M235T) and coronary heart disease. *Jiangsu Med J* 2001; 27: 254-6.

47. Zhu T, Xu Z, Yang Z, Xie Y, Zhang Z, Jia Y, et al. The association of polymorphism of three genes of the renninangiotensin system with coronary heart disease. *Jiangsu Med J* 2002; 28: 745-6.
48. Zhu Y, Hs Long H. Relationship of angiotensinogen gene polymorphism and myocardial infarction in Chinese. *Chin J Cardiol* 2002; 12: 67.
49. Gu W, Zhu J, Pan M, Yuan Z, Gong Y, Cui Z, et al. Effects of angiotensinogen and apolipoprotein E gene polymorphisms on coronary heart disease. *J NanTong Med Univ* 2003; 4: 394-5.
50. Li L FXH, Zhang YW. Association between angiotensinogen gene polymorphism and severity of coronary artery stenosis. *J Med Forum* 2004; 25: 12-4.
51. Zhu T, Pan J, Shen Z, Zhao Y, Shen Z. Gene polymorphisms of reninangiotensin system and coronary thrombosis disease. *Journal of Experimental Hematology* 2004; 12: 674-9.
52. Ren J, Jia Y, Lv J, Guo X. Relationship between angiotensinogen gene M235T polymorphism and coronary heart disease. *Journal of Shanxi Medical University* 2005; 36: 439-42.
53. Liang Q DYG, Yang XL. Research of ACE, angiotensinogen and endothelial nitric oxide synthase (eNOS) gene polymorphisms in predisposition to CHD with chip technology. *Hebei Medicine* 2006; 12: 403-6.
54. Tsai CT, Hwang JJ, Ritchie MD, Moore JH, Chiang FT, Lai LP, et al. Renin-angiotensin system gene polymorphisms and coronary artery disease in a large angiographic cohort: detection of high order gene-gene interaction. *Atherosclerosis* 2007; 195: 172-80.
55. Niu Y, Luo L, Peng J, Mei X, Peng Z, Gong W. Relationship between Renin-Angiotensin System Gene Polymorphism and Ischemic Chronic Heart Failure. *Academic Journal of Sun Yat-sen University of Medi* 2008; 29: 168-72.
56. Zhu YG, Wang FJ, Fu ZQ, Xiang YD. Association between the gene polymorphism of angiotensinogen-converting enzyme and coronary artery disease. *Chin J Arterioscler* 2010; 18: 405-8.
57. Peng J, Luo LY, Li T, Mei X, Niu YQ, Gong WX. Relationship between renninangiotensin system gene polymorphism and ischemic chronic heart failure in aged coronary artery disease patient. *Chinese Circulation Journal* 2011; 26: 93-6.
58. Niu T, Chen X, Xu X. Angiotensin converting enzyme gene insertion/deletion polymorphism and cardiovascular disease: therapeutic implications. *Drugs* 2002; 62: 977-93.
59. Lanz JR, Pereira AC, Lemos PA, Martinez E, Krieger JE. Angiotensinogen M235T polymorphism is associated with coronary artery disease severity. *Clin Chim Acta* 2005; 362: 176-81.
60. Jeunemaitre X, Soubrier F, Kotelevtsev YV, Lifton RP, Williams CS, Charru A, et al. Molecular basis of human hypertension: role of angiotensinogen. *Cell* 1992; 71: 169-80.
61. Horiuchi M, Akishita M, Dzau VJ. Recent progress in angiotensin II type 2 receptor research in the cardiovascular system. *Hypertension* 1999; 33: 613-21.
62. Henrion D, Kubis N, Lévy BI. Physiological and pathophysiological functions of the AT(2) subtype receptor of angiotensin II: from large arteries to the microcirculation. *Hypertension* 2001; 38: 1150-7.
63. Dietz R, von Harsdorf R, Gross M, Krämer J, Gulba D, Willenbrock R, et al. Angiotensin II and coronary artery disease, congestive heart failure, and sudden cardiac death. *Basic Res Cardiol* 1998; 93 Suppl 2: 101-8.
64. Keidar S, Attias J, Heinrich R, Coleman R, Aviram M. Angiotensin II atherogenicity in apolipoprotein E deficient mice is associated with increased cellular cholesterol biosynthesis. *Atherosclerosis* 1999; 146: 249-57.
65. Li W, Li J, Hao P, Chen W, Meng X, Li H, et al. Imbalance between angiotensin II and angiotensin-(1-7) in human coronary atherosclerosis. *J Renin Angiotensin Aldosterone Syst* 2016; 17.
66. Katsuya T, Koike G, Yee TW, Sharpe N, Jackson R, Norton R, et al. Association of angiotensinogen gene T235 variant with increased risk of coronary heart disease. *Lancet* 1995; 345: 1600-3.
67. Liang X, Qiu J, Liu X, Li X, Zhao S, Wang J, et al. Polymorphism of angiotensinogen gene M235T in myocardial infarction and brain infarction: a meta-analysis. *Gene* 2013; 529: 73-9.
68. Walker AR, Walker BF, Segal I. Some puzzling situations in the onset, occurrence and future of coronary heart disease in developed and developing populations, particularly such in sub-Saharan Africa. *J R Soc Promot Health* 2004; 124: 40-6.
69. Micha R, Peñalvo JL, Cudhea F, Imamura F, Rehm CD, Mozaffarian D. Association Between Dietary Factors and Mortality From Heart Disease, Stroke, and Type 2 Diabetes in the United States. *JAMA* 2017; 317: 912-24.
70. Poelman M, Strak M, Schmitz O, Hoek G, Karssenberg D, Helbich M, et al. Relations between the residential fast-food environment and the individual risk of cardiovascular diseases in The Netherlands: A nationwide follow-up study. *Eur J Prev Cardiol* 2018; 25: 1397-405.